arXiv:1602.02745v2 [math.CV] 12 Feb 2016

On measures which generate the scalar product in a space of rational functions.

Victor Katsnelson

Abstract. Let $z_1, z_2, \dots, z_n$ be pairwise different points of the unit disc and $\mathcal{L}(z_1, z_2, \dots z_n)$ be the linear space generated by the rational fractions $\frac{1}{t-z_1}, \frac{1}{t-z_2}, \cdots, \frac{1}{t-z_n}$. Every non-negative measure σ on the unit circle $\mathbb{T}$ generates the scalar product

$$\langle f\,,\, g\rangle_{L^2_\sigma} = \int\limits_{\mathbb{T}} f(t)\,\overline{g(t)}\,\sigma(dt), \quad \forall\, f, g \,\in\, L^2_\sigma.$$

The measures σ are described which satisfy the condition

$$\langle f\,,\, g\rangle_{L^2_\sigma} = \langle f\,,\, g\rangle_{L^2_m}, \quad \forall\, f, g \in \mathcal{L}(z_1, z_2, \dots z_n),$$

where m is the normalized Lebesgue measure on $\mathbb{T}$.

Mathematics Subject Classification (2010). 46E20.

Keywords. Non-negative measures on the unite circle; rational fractions with poles in the unit disc.

- $\mathbb{R}$ stands for the set of all real numbers.
- $\mathbb{C}$ stands for the set of all complex numbers.
- $\mathbb{T}$ is the unite circle: $\mathbb{T} = \{t \in \mathbb{C} : |t| = 1\}$.
- $\mathbb{D}_+$ is the open unite disc: $\mathbb{D}_+ = \{z \in \mathbb{C} : |z| < 1\}$.
- $\mathbb{D}_-$ is the exterior of the unite circle: $\mathbb{D}_- = \{z \in \mathbb{C} : 1 < |z| \leq \infty\}$.

The main objects of this note are spaces of functions on the unit circle $\mathbb{T}$ and measures generated scalar products in such spaces. Let us set some notation.

For a non-negative measure σ on $\mathbb{T}$, the space L^2_σ is the set of all square integrable functions with respect to the measure σ. The space L^2_σ is provided by natural linear operations and by the scalar product

$$\langle f\,,\, g\rangle_{L^2_\sigma} = \int\limits_{\mathbb{T}} f(t)\,\overline{g(t)}\,\sigma(dt). \quad \forall\, f, g \,\in\, L^2_\sigma.$$

Let $z_1, z_2, \dots, z_n$, $n < \infty$, be a sequence of pairwise different points of the unit disc $\mathbb{D}_+$:

$$z_k \in \mathbb{D}_+, \quad 1 \le k \le n, \tag{1a}$$

$$z_p \ne z_q, \quad 1 \le p, q \le n. \tag{1b}$$

We relate two objects to this sequence. The first object is the Blaschke product[1]

$$B(t) = \prod_{1 \le k \le n} \frac{z_k - t}{1 - \overline{z_k} t} \cdot \frac{|z_k|}{z_k}. \tag{2}$$

The second object is the sequence

$$e(t, z_k) = \frac{1}{t - z_k}, \quad 1 \le k \le n, \tag{3}$$

of rational fractions.

Let $\mathcal{L}(z_1, z_2, \dots, z_n)$ be the linear space generated by the functions $e(t, z_k)$, $1 \le k \le n$. In other words, $\mathcal{L}(z_1, z_2, \dots, z_n)$ is the set of all linear combinations $\sum\limits_{1 \le k \le h} \alpha_k \, e(t, z_k)$, where α_k, $1 \le k \le n$, are arbitrary complex numbers.

Functions which belong to the linear space $\mathcal{L}(z_1, z_2, \dots, z_n)$ are continuous on the unit circle $\mathbb{T}$. Therefore they belong to every space L^2_σ, where σ is a nonnegative measure on $\mathbb{T}$. If

$$f(t) = \sum_{1 \le k \le n} \xi_k \, e(t, z_k), \quad g(t) = \sum_{1 \le k \le n} \eta_k \, e(t, z_k) \tag{4}$$

belong to $\mathcal{L}(z_1, z_2, \dots, z_n)$, then their scalar product in the space L^2_σ is equal to

$$\langle f \, , \, g \rangle_{L^2_\sigma} = \sum_{1 \le k, l \le n} \xi_k \, \overline{\eta_l} \, \langle e(., z_k) \, , \, e(., z_l) \rangle_{L^2_\sigma}. \tag{5}$$

Among all nonnegative measures σ on $\mathbb{T}$, we distinguish the normalised Lebesgue measure $m(dt)$.

We discuss the following

Problem 1. *Let pairwise different points $z_1, z_2, \dots, z_n$ from $\mathbb{D}_+$ be given. How to describe those nonnegative measures σ on $\mathbb{T}$ for which the property*

$$\langle f \, , \, g \rangle_{L^2_\sigma} = \langle f \, , \, g \rangle_{L^2_m}, \quad \forall \, f, g \in \mathcal{L}(z_1, z_2, \dots, z_n). \tag{6}$$

holds?

Definition 1. Given the set $z_1, z_2, \dots, z_n$ of pairwise different points from $\mathbb{D}_+$, let $\Sigma(z_1, z_2, \dots, z_n)$ be the set of all non-negative measures on $\mathbb{T}$ for which the property (6) holds.

Lemma 2. *Given the set $z_1, z_2, \dots, z_n$ of pairwise different points from $\mathbb{D}_+$, then the set $\Sigma(z_1, z_2, \dots, z_n)$ is not empty, convex, and bounded set of measures which is closed with respect to the weak topology.*

[1] If $z_k = 0$ for some k, then the appropriate factor of the Blaschke product (2) is equal just to t, but not to $\frac{z_k - t}{1 - \overline{z_k} t} \cdot \frac{|z_k|}{z_k}$.

Proof. The set $\Sigma(z_1, z_2, \dots, z_n)$ is not empty: the Lebesgue measure m belongs to this set. The convexity of this set and its weak closeness are evident. The $\Sigma(z_1, z_2, \dots, z_n)$ is bounded. In Lemma 13 we establish the estimate

$$\sigma(\mathbb{T}) \le \frac{1 + B(0)}{1 - B(0)}, \quad \forall\, \sigma \in \Sigma(z_1, z_2, \dots, z_n). \tag{7}$$

(We emphasize that the inequalities (36) holds.) □

In view of (5), the equality (6) is equivalent to the totality of equalities

$$\langle\, e(., z_k)\,,\, e(., z_l)\,\rangle_{L^2_\sigma} = \langle\, e(., z_k)\,,\, e(., z_l)\,\rangle_{L^2_m}, \quad 1 \le k, l \le n. \tag{8}$$

The expressions in the right hand side of (8) can be calculated explicitly:

$$\langle\, e(., z_k)\,,\, e(., z_l)\,\rangle_{L^2_m} = \frac{1}{1 - z_k\overline{z_l}}, \quad 1 \le k, l \le n. \tag{9}$$

The expressions in the left hand side of (8) can be presented as

$$\langle\, e(., z_k)\,,\, e(., z_l)\,\rangle_{L^2_\sigma} = \int\limits_{\mathbb{T}} \frac{1}{t - z_k} \frac{1}{\overline{t - z_l}}\, \sigma(dt), \quad 1 \le k, l \le n. \tag{10}$$

Thus the Problem 1 can be reformulated as follows:

Problem 1′. *Let pairwise different points $z_1, z_2, \dots, z_n$ from $\mathbb{D}_+$ be given. How to describe such nonnegative measures σ on $\mathbb{T}$ for which the totality of equalities*

$$\int\limits_{\mathbb{T}} \frac{1}{t - z_k} \frac{1}{\overline{t - z_l}}\, \sigma(dt) = \frac{1}{1 - z_k\overline{z_l}}, \quad 1 \le k, l \le n, \tag{11}$$

hold?

It turns out that the Problem 1′ can be reduced to a very special Newanlinna-Pick interpolation problem.

Let σ be a nonnegative measure on $\mathbb{T}$. We associate the function

$$\varphi_\sigma(z) = \int\limits_{\mathbb{T}} \frac{t + z}{t - z}\, \sigma(dt), \quad z \in \mathbb{D}_+, \tag{12}$$

with this σ. It is clear that the function φ_σ is holomorphic in $\mathbb{D}_+$.

If $\zeta', \zeta'' \in \mathbb{D}_+$, then

$$\frac{\varphi_\sigma(\zeta') + \overline{\varphi_\sigma(\zeta'')}}{2(1 - \zeta'\overline{\zeta''})} = \int\limits_{\mathbb{T}} \frac{1}{t - \zeta'}\, \frac{1}{\overline{t - \zeta''}}\, \sigma(dt). \tag{13}$$

For $\zeta' = z_k$, $\zeta'' = z_l$, the expression in the right hand side of (13) coincides with the expression in the left hand side of (11). Thus the equalities (11) can be rewritten in the form

$$\frac{\varphi_\sigma(z_k) + \overline{\varphi_\sigma(z_l)}}{1 - z_k\overline{z_l}} = \frac{2}{1 - z_k\overline{z_l}}, \quad 1 \le k, l \le n,$$

that is

$$\varphi_\sigma(z_k) + \overline{\varphi_\sigma(z_l)} = 2, \quad 1 \le k, l \le n. \tag{14}$$

Thus the following statement is proved.

Lemma 3. *Let $z_1, z_2, \dots, z_n$ be pairwise different points of the unit disc $\mathbb{D}_+$. For a non-negative measure σ on $\mathbb{T}$, let φ_σ be the function associated with the measure σ according to* (12).

1. *If $\sigma \in \Sigma(z_1, z_2, \dots, z_n)$, then the equalities* (14) *hold.*
2. *If the equalities* (14) *hold, then the $\sigma \in \Sigma(z_1, z_2, \dots, z_n)$.*

We consider the equalities (14) as a system of equations with respect to the values $\varphi_\sigma(z_k)$.

Lemma 4. *Let $\varphi_1, \varphi_2, \dots, \varphi_n$ be complex numbers satisfying the system of equations*

$$\varphi_k + \overline{\varphi_l} = 2, \quad 1 \le k, l \le n. \tag{15}$$

Then there exists a real number β such that

$$\varphi_k = 1 - i\beta, \quad 1 \le k \le n. \tag{16}$$

Vice versa, if β is an arbitrary real number and φ_k are defined by (16)*, then the equalities* (15) *hold.*

Thus the set of solutions of the system of equations (15) *is an one-parametric family. The arbitrary real number β is a free parameter which parameterizes this family according to* (16).

Proof. Let $\boldsymbol{e} = [1, 1, \dots, 1]$ and be $\boldsymbol{\varphi} = [\varphi_1, \varphi_2, \dots, \varphi_n]$ be $1 \times n$ matrices (rows), $\boldsymbol{e}^*$ and $\boldsymbol{\varphi}^*$ be the Hermitian conjugate matrices (columns). The system (15) can be rewritten in a matrix form

$$\boldsymbol{\varphi}^* \boldsymbol{e} + \boldsymbol{e}^* \boldsymbol{\varphi} = 2\, \boldsymbol{e}^* \boldsymbol{e}\,. \tag{17}$$

The matrix in the right hand side of (17) is of rank one. Thus $\boldsymbol{\varphi}$ must be of the form $\boldsymbol{\varphi} = \zeta\, \boldsymbol{e}$, where $\zeta \in \mathbb{C}$. (Otherwise the matrix in the left hand side of (17) is of rank two.) Substituting $\boldsymbol{\varphi} = \zeta\, \boldsymbol{e}$ into (17), we see that $\boldsymbol{\varphi} = (1 - i\beta)\, \boldsymbol{e}$, where $\beta \in \mathbb{R}$. □

From the equalities (14) and Lemma 4 it follows that there exists a real number β such that the equalities

$$\varphi_\sigma(z_k) = 1 - i\beta, \quad 1 \le k \le n. \tag{18}$$

hold.

Given a non-negative measure σ on $\mathbb{T}$ and a real number β, we associate the function

$$c_{\sigma,\beta}(z) = i\beta + \int_{\mathbb{T}} \frac{t+z}{t-z}\, \sigma(dt), \quad z \in \mathbb{D}_+. \tag{19}$$

with these σ and β. It is clear that

$$\operatorname{Re} c_{\sigma,\beta}(z) = h_\sigma(z), \tag{20}$$

where

$$h_\sigma(z) = \int_{\mathbb{T}} \frac{1 - |z|^2}{|t-z|^2}\, \sigma(dt), \quad z \in \mathbb{D}_+. \tag{21}$$

The equalities (18) can be rewritten as

$$c_{\sigma,\beta}(z_k) = 1, \quad 1 \le k \le n. \tag{22}$$

Lemma 3 can be reformulated as follows.

Lemma 5. *Let $z_1, z_2, \dots, z_n$ be pairwise different numbers from $\mathbb{D}_+$. For a non-negative measure σ on $\mathbb{T}$ and a real number β, let $c_{\sigma,\beta}(z)$ be the function associated with these σ and β according to* (12).

1. *If $\sigma \in \Sigma(z_1, z_2, \dots, z_n)$, then there exists the real number β such that the interpolation conditions* (22) *hold for the function $c_{\sigma,\beta}$.*
2. *If the interpolation conditions* (22) *hold for some function $c_{\sigma,\beta}$, then $\sigma \in \Sigma(z_1, z_2, \dots, z_n)$.*

The functions of the form (22), where σ is an arbitrary non-negative measure on $\mathbb{T}$ and β is an arbitrary real number, can be characterized by their properties.

Definition 6. *The Caratheodory class $\boldsymbol{\mathcal{C}}$* is the class of all functions $c(z)$ which possess the properties:

1. The function $c(.)$ is defined and holomorphic in the disc $\mathbb{D}_+$.
2. The real part of the function $c(.)$ is non-negative in $\mathbb{D}$:

$$\operatorname{Re} c(z) \ge 0 \quad \forall z \in \mathbb{D}_+. \tag{23}$$

The Representation Theorem for the Caratheodory class.

1. *Let σ be a non-negative measure on $\mathbb{T}$, β is a real number, and $c_{\sigma,\beta}(.)$ be the function defined by the equality* (19). *Then the function $c_{\sigma,\beta}(.)$ belongs to the Caratheodory class $\boldsymbol{\mathcal{C}}$.*
2. *Let $c(.)$ be a function which belongs to the Caratheodory class $\boldsymbol{\mathcal{C}}$. Then the function $c(.)$ is representable in the form* (19)*:*

$$c(z) = c_{\sigma,\beta}(z) \tag{24}$$

with some non-negative measure σ and real number β. The measure σ and number β are determined by the function $c(.)$ uniquely.

The next statement is an immediate consequence of Lemma 5 and of the Representation Theorem.

Lemma 7. *Let $z_1, z_2, \dots, z_n$ be pairwise different points from $\mathbb{D}_+$. Then the set of non-negative measures σ belonging to the set $\Sigma(z_1, z_2, \dots, z_n)$ coincides with the set of measures σ appearing as the representing measures,* (24)*, for those functions $c(.) \in \boldsymbol{\mathcal{C}}$ which satisfy the interpolating conditions*

$$c(z_k) = 1, \quad 1 \le k \le n. \tag{25}$$

Thus the original problem, i.e. Problem 1, is related to the following interpolation problem in the Caratheodory class.

Problem $\mathbf{2}_{\boldsymbol{\mathcal{C}}}$. *Let $z_1, z_2, \dots, z_n$ be pairwise different points from $\mathbb{D}_+$.* The function c is a solution of the Problem $\mathbf{2}_{\boldsymbol{\mathcal{C}}}$ *if $c \in \boldsymbol{\mathcal{C}}$, and the interpolation conditions* (25) *are satisfied.*

This interpolation problem is the *Nevanlinna-Pick problem in the class* $\mathcal{C}$ with generic interpolation nodes, but with very special interpolation values.

There is the very well developed machinery for study the Nevanlinna-Pick problems in various functional classes. (See [Kov].) This machinery includes the solvability criteria as well as the description of the set of all solutions in the case of solvability. We do not need to use such machinery. The set of solutions of the Nevanlinna-Pick problem (25) in the class $\boldsymbol{\mathcal{C}}$ can be easily describe without using the mentioned general theory.

Definition 8. *The Schur class* $\boldsymbol{\mathcal{S}}$ is the class of all functions $s(z)$ which possess the properties:

1. The function $s(.)$ is defined and holomorphic in the disc $\mathbb{D}_+$.
2. The absolute value of the function $s(.)$ does not exceed one in $\mathbb{D}$:

$$|s(z)| \leq 1 \quad \forall z \in \mathbb{D}_+. \tag{26}$$

The fractional linear transformation

$$c(z) = \frac{1 + s(z)}{1 - s(z)} \tag{27}$$

establishes one-to one correspondence between classes $\boldsymbol{\mathcal{C}}$ and $\boldsymbol{\mathcal{S}}$.

It is clear that if the functions c and s are related by the transformation (27), then the interpolation conditions

$$s(z_k) = 0, \quad 1 \leq k \leq n, \tag{28}$$

for the function s correspond to the interpolation conditions (25) to the function c.

Thus the interpolation Problem $2_{\mathcal{C}}$ in the class $\boldsymbol{\mathcal{C}}$ is reduced to the following interpolation problem in the class $\boldsymbol{\mathcal{S}}$.

Problem $2_{\boldsymbol{\mathcal{S}}}$. *Let $z_1, z_2, \ldots, z_n$ be pairwise different points from $\mathbb{D}_+$. The function s is a solution of the Problem $\mathbf{2}_{\boldsymbol{\mathcal{S}}}$ if $s \in \boldsymbol{\mathcal{S}}$, and the interpolation conditions* (28) *are satisfied.*

Lemma 9. *Let c and s be functions related by the fractional linear transformation* (27)*. The function c is a solution of the interpolation Problem $2_{\mathcal{C}}$ if and only if the function s is a solution of the interpolation Problem $2_{\mathcal{S}}$.*

Lemma 10. *The interpolation* Problem $2_{\boldsymbol{\mathcal{S}}}$ *is solvable. The formula*

$$s(z) = B(z)\,\omega(z), \tag{29}$$

where $B(z)$ is the Blaschke product (2) *constructed from the given interpolation nodes z_k, $1 \leq k \leq n$, parameterizes the set of all solutions s of the* Problem $2_{\boldsymbol{\mathcal{S}}}$ *by means of arbitrary functions $\omega \in \boldsymbol{\mathcal{S}}$:*

1. *If s is a solution of the* Problem $2_{\boldsymbol{\mathcal{S}}}$*, then s is representable in the form* (29)*, where $\omega \in \boldsymbol{\mathcal{S}}$.*
2. *If $\omega \in \boldsymbol{\mathcal{S}}$ and s is related to ω by* (29)*, then s is a solution of* Problem $2_{\boldsymbol{\mathcal{S}}}$.

3. *The correspondence* (29) *between parameters* $\omega \in \boldsymbol{\mathcal{S}}$ *and solutions* s *of* Problem $2_{\boldsymbol{\mathcal{S}}}$ *is one-to-one.*

In view of the mentioned relation between Problems $2_{\mathcal{S}}$ and $2_{\mathcal{C}}$, the following result is established.

Lemma 11. *The interpolation* Problem $2_{\boldsymbol{\mathcal{C}}}$ *is solvable. The formula*

$$c(z) = \frac{1 + B(z)\,\omega(z)}{1 - B(z)\,\omega(z)}, \tag{30}$$

where $B(z)$ *is the Blaschke product* (2) *constructed from the given interpolation nodes* z_k, $1 \le k \le n$, *parameterizes the set of all solutions* c *of the* Problem $2_{\boldsymbol{\mathcal{C}}}$ *by means of arbitrary functions* $\omega \in \boldsymbol{\mathcal{S}}$*:*

1. *If* c *is a solution of the* Problem $2_{\boldsymbol{\mathcal{C}}}$*, then* c *is representable in the form* (30)*, where* $\omega \in \boldsymbol{\mathcal{S}}$*.*
2. *If* $\omega \in \boldsymbol{\mathcal{S}}$ *and* c *is related to* ω *by* (30)*, then* c *is a solution of* Problem $2_{\boldsymbol{\mathcal{C}}}$*.*
3. *The correspondence* (30) *between parameters* $\omega \in \boldsymbol{\mathcal{S}}$ *and solutions* c *of* Problem $2_{\boldsymbol{\mathcal{C}}}$ *is one-to-one.*

Lemmas 5 and 11 allow to describe the set of all measures σ which belongs to the set $\Sigma(z_1, z_2, \dots, z_n)$. Holomorphic functions $c_{\sigma,\beta}$ from the Caratheodory class appear in such description. However it is more convenient to formulate the final result not in terms of functions of the Caratheodory class but in terms of real parts such functions. This allows to eliminate the value β which is not related to the original problem.

Definition 12. *The Herglotz class* $\boldsymbol{\mathcal{H}}$ is the class of all functions $h(z)$ which possess the properties:

1. The function $h(.)$ is defined and harmonic in the disc $\mathbb{D}_+$.
2. The function $h(.)$ is non-negative in $\mathbb{D}$:

$$h(z) \ge 0, \quad \forall z \in \mathbb{D}_+. \tag{31}$$

The following theorem characterizes functions $h \in \boldsymbol{\mathcal{H}}$ as functions which admit the representation

$$h(z) = \int_{\mathbb{T}} \frac{1 - |z|^2}{|t - z|^2}\,\sigma(dt), \quad z \in \mathbb{D}_+, \tag{32}$$

with a non-negative measure σ.

Riesz-Herglotz Representation Theorem.

1. *Let* h *be a function from the class* $\boldsymbol{\mathcal{H}}$*. Then the function* h *is representable in the form* (32)*, where* σ *is a non-negative measure on* $\mathbb{T}$*.*
2. *Let* σ *be a non-negative measure on* $\mathbb{T}$ *and the function* h *be constructed from this* σ *according to* (32)*. Then* $h \in \boldsymbol{\mathcal{H}}$*.*
3. *The correspondence* (32) *between* h *and* σ *is one-to-one.*

The following Theorem is the main result of this note.

Theorem. *Let $z_1, z_2, \dots, , z_n$ be pairwise different points from $\mathbb{D}_+$ and $B(z)$ be the Blaschke product constructed from these points,* (2).
The formula

$$h(z) = \operatorname{Re} \frac{1 + B(z)\,\omega(z)}{1 - B(z)\,\omega(z)}, \quad z \in \mathbb{D}_+, \tag{33}$$

parameterises the set $\Sigma(z_1, z_2, \dots, z_n)$. The function $\omega \in \boldsymbol{\mathcal{S}}$ serves as a parameter.

1. *Let $\omega(z)$ be a function from the Schur class $\boldsymbol{\mathcal{S}}$ and the function h be constructed from this ω by the formula* (33).
   Then
   (a) *The function h is a harmonic function from the Herglotz class $\boldsymbol{\mathcal{H}}$.*
   (b) *The measure σ which appears in the Riesz-Herglotz representation* (32) *of the function h belongs to the set $\Sigma(z_1, z_2, \dots, z_n)$.*
2. *Let σ be a measure which belongs to the set $\Sigma(z_1, z_2, \dots, z_n)$. Let h be the function which is constructed from this σ according to* (32).
   Then there exists the function $\omega(z)$ from the Schur class $\boldsymbol{\mathcal{S}}$ such that the equality (33) *holds.*
3. *Different functions h correspond to different parameters ω.*

The parametrization (33) allows to estimate the values $\sigma(\mathbb{T})$ for the measures $\sigma \in \Sigma(z_1, z_2, \dots, z_n)$.

Lemma 13. *Let $z_1, z_2, \dots, z_n$ be pairwise different points from $\mathbb{D}_+$. Then*

$$\max\{\sigma(\mathbb{T}) : \sigma \in \Sigma(z_1, z_2, \dots, z_n)\} = \frac{1 + B(0)}{1 - B(0)}, \tag{34a}$$

$$\min\{\sigma(\mathbb{T}) : \sigma \in \Sigma(z_1, z_2, \dots, z_n)\} = \frac{1 - B(0)}{1 + B(0)}. \tag{34b}$$

Proof. Let σ be the measure which represents the function h from (33). Then

$$\sigma(\mathbb{T}) = h(0) = \operatorname{Re} \frac{1 + B(0)\omega(0)}{1 - B(0)\omega(0)}. \tag{35}$$

Since $B(0) = \prod\limits_{1 \le k \le n} |z_k|$, the inequalities

$$0 \le B(0) < 1 \tag{36}$$

hold. When the parameter ω runs over the whole Schur class $\boldsymbol{\mathcal{S}}$, the value $\omega(0)$ runs over the *closed* unit disc $\overline{\mathbb{D}}_+ = \mathbb{D}_+ \cup \mathbb{T}$. Therefore the value in the right hand side of (35) attained its maximum for $\omega(0) = +1$ and its minimum for $\omega(0) = -1$. The maximum corresponds to the choice of the parameters $\omega(z) \equiv +1$, the minimum corresponds to the choice of the parameters $\omega(z) \equiv -1$. □

Remark. The space $\mathcal{L}(z_1, z_2, \dots, z_n)$ can be identified with the subspace $H^2_- \ominus B^{-1}H^2_-$ of the Hilbert space L^2_m, where H^2_- is the Hardy class in the domain $\mathbb{D}_-$ and $\ominus$ is the orthogonal complement with respect to the scalar product in L^2_m.

References

[Kov] Kovalishina, I.V. *Analytic Theory of a class of interpolation problem.* Mathematics of the USSR-Izvestiya, 1984, **22**:3, 419-463.

Victor Katsnelson
Department of Mathematics
The Weizmann Institute
76100, Rehovot
Israel
e-mail: victor.katsnelson@weizmann.ac.il; victorkatsnelson@gmail.com